

Patient Name: \_\_\_\_\_

Date Prepared: \_\_\_\_\_

### Compounding Formulation Record

Rx: \_\_\_\_\_

Recipe ID#: \_\_\_\_\_ (IF AVAILABLE)

Batch #: \_\_\_\_\_ Beyond Use Date: 30 Days Good For Use Until: \_\_\_\_/\_\_\_\_/\_\_\_\_

#### Compounding Formulation Record for: FIRST®-Metronidazole 50 mg/mL Suspension

**WARNING**  
Metronidazole has been shown to be carcinogenic in mice and rats. Unnecessary use of the drug should be avoided.  
See PRECAUTIONS and usage described in approved labeling for metronidazole containing products for additional information.

CLASS: RX      DOSAGE FORM: Suspension      ROUTE OF ADMINISTRATION: Oral

Ingredient Name	Dispensed Quantity	NDC/UPC #	Manufacturer	Lot #	Exp. Date of Ingredient	Prepared By:	Verified By:
Metronidazole Benzoate Powder		65628-202-05	Azurity				
Diluent (FIRST®- Grape II Suspension)		65628-202-05	Azurity				

**Prior to compounding, store FIRST®-Metronidazole Compounding Kit at room temperature 15° - 30°C (59° - 86°F)  
Store the final compounded formulation at room temperature 15° - 30°C (59° - 86°F) for up to 30 days.**

**FIRST®-Metronidazole 5 FL OZ (150 mL) Kit (NDC 65628-202-05) contains 12 g Metronidazole Benzoate Powder + 141 mL Diluent (FIRST®-Grape II Suspension)**

Total Quantity	mL						
----------------	----	--	--	--	--	--	--

**SEE FULL PACKAGE INSERT FOR ADDITIONAL INFORMATION. THE RECORD DOES NOT INCLUDE ALL THE INFORMATION NEEDED TO USE THE PRODUCT SAFELY AND EFFECTIVELY.**

**COMPOUNDING INSTRUCTIONS:**

- FIRST®-Metronidazole 50 Compounding Kit contains pre-measured metronidazole benzoate powder and Diluent (FIRST®- Grape II Suspension).
- Important - Hold the neck of the bottle containing metronidazole benzoate powder and tap the bottom edges on a hard surface to loosen the powder. Remove the cap from the bottle. Tap the top of the induction seal liner to loosen any powder which may have adhered to the liner. Carefully and slowly peel back the inner foil seal liner from the bottle.
- Shake the Diluent bottle for a few seconds. Open the suspension bottle and pour approximately half of the Diluent into the metronidazole benzoate powder bottle.
- Important:** Replace the cap and shake the metronidazole benzoate powder bottle ***well for approximately 60 seconds*** to ensure a well-mixed suspension.
- Empty the remaining Diluent ***into the metronidazole benzoate powder bottle, and allow the remaining Diluent to drain into the powder bottle for 10 seconds.***
- Replace the cap and shake the metronidazole benzoate powder bottle ***Well again for approximately 60 seconds.***
- Dispense the metronidazole benzoate oral suspension in the bar coded bottle to the patient.
- Important:** Be sure to instruct your patient to wait for at least one hour before administering the first dose and also to ***shake the suspension Well*** before each use in order to ensure a well- mixed suspension.

**WARNINGS:**

- Central and Peripheral Nervous System Effects

**AUXILIARY LABELS:**

- For oral use only
- Avoid contact with eyes
- Keep container tightly closed
- Keep out of the reach of children
- Protect from light
- Protect from freezing
- When stored at room temperature the beyond-use date of the compounded product is ***no later than 30 days***

**REFERENCE:**

FIRST® - METRONIDAZOLE 50 MG/ML [PACKAGE INSERT].  
 Wilmington, MA: Azurity Pharmaceuticals, Inc. 2020  
**PART CMP-687 | REV. 00**

